

Umbilical epidermoid cyst: an unusual cause of umbilical symptoms

James H. McClenathan, MD

In addition to the most likely diagnoses for umbilical signs in adults (i.e., umbilical hernia, metastatic tumour, endometriosis, congenital abnormalities and gallstones retained or spilled at the umbilicus), the differential diagnosis should include epidermoid cyst arising from the skin of the umbilicus. Indeed, to determine the most appropriate treatment for patients with umbilical signs or symptoms, physicians should be aware that deep-seated epidermoid cysts may occur at the umbilicus and that these cysts may extend below the midline fascia. On the basis of our observations reported here and those reported elsewhere,¹⁻⁴ at our institution, we believe that these cysts must be completely excised. We have not encountered any recurrence with this treatment. This communication describes 7 patients with symptoms resulting from an epidermoid cyst of the umbilicus.

Patients and treatment

Between 1984 and 2001, 7 patients (2 male, 5 female), ranging in age from 9 to 57 years, were treated at our facility for brief or long-standing umbilical signs or symptoms resulting from an epidermoid cyst (Table 1). Five of these patients had umbilical pain, which in several patients radiated away from the umbilicus. Four patients had a palpable mass; 4 had umbilical redness. Two patients had umbilical drainage that was described as having the odour of sebum. Use of cotton swabs and good lighting sometimes enabled us to see a punctum or to extract sebaceous material from the depths of the umbilicus.

Six patients were treated surgically. Where possible, the cyst was completely excised with a small piece of skin attached. This excision was more difficult in patients with active infection. In 2

patients, the major component of the epidermoid cyst was located below the fascia and communicated with the skin through a tiny fascial opening. In 1 patient, thrombophlebitis of a superficial inferior epigastric vein developed in the right lower quadrant secondary to infection.

Discussion

In most patients having umbilical symptoms who seek medical attention, an umbilical hernia is diagnosed. We have previously reported our experience treating pilonidal sinus disease of the umbilicus¹ and have treated patients with suture granuloma, endometriosis, and urachal and omphalomesenteric anomalies. In a review of umbilical inflammatory conditions, Goldberg and associates² classified these conditions as primary or secondary and also discussed urachal and

Table 1

Characteristics of 7 Patients Treated for Umbilical Signs or Symptoms Caused by Epidermoid Cyst, 1984–2001

Year	Age, yr	Sex	Duration of signs/symptoms	Pain	Mass	Redness	Drainage	Location	Treatment
1984	30	M	n/a	Yes	No	Yes	n/a	Below fascia	Excision
1993	52	F	Years	No	Yes	No	No	Above fascia	Excision
1994	39	F	1 d	Yes	Yes	Yes	No	Above fascia	Excision
1997	30	F	Months	Yes	Yes	Yes	Yes	Below fascia	Excision
2000	57	F	Weeks	Yes	No	No	Yes	Above fascia	Antibiotics
2000	51	M	2 wk	Yes	No	Yes	No	Above fascia	Excision
2001	9	F	1 yr	No	Yes	No	No	Above fascia	Excision

n/a = not available.

From the Department of Surgery, Kaiser Permanente Medical Center, Santa Clara, Calif., and Clinical Professor of Surgery, Stanford University School of Medicine, Stanford, Calif.

Accepted for publication Aug. 21, 2001.

Correspondence to: Dr. James H. McClenathan, Department of Surgery, Kaiser Permanente Santa Clara Medical Center, 900 Kiely Blvd., Santa Clara, CA 95051 USA; fax 408 236-4868; james.mcclenathan@kp.org

omphalomesenteric abnormalities, umbilical polyps, primary omphalitis, pilonidal sinus, primary umbilical carcinoma and inflammation secondary to intra-abdominal or retroperitoneal infection. However, they did not mention infected epidermoid cyst in their list of causes of umbilical inflammation. Molderez and colleagues³ described a series of 22 patients with umbilical discharge. Among 17 patients who had umbilical discharge of previously undetermined cause, discharge in 10 was found to result from a condition of embryologic origin. In 1 patient, umbilical discharge was caused by a sebaceous cyst.³ Cullen's textbook of diseases of the umbilicus and urachus,⁴

published in 1916, described walnut-sized dermoid (or atheromatous) cysts of the umbilicus reported in only 5 patients. This text stated the following: these cysts originate from the umbilicus; they contain sebaceous material that yields epithelial cells, fat droplets and cholesterol crystals; they are lined by squamous epithelium; and they do not contain hair. We recommend complete excision of epidermoid cysts. We have not seen recurrence with that treatment.

Acknowledgements: The Medical Editing Department, Kaiser Foundation Research Institute, provided editorial assistance.

References

1. McClenathan JH. Umbilical pilonidal sinus. *Can J Surg* 2000;43(3):225.
2. Goldberg R, Pritchard B, Gelbard M. Umbilical inflammatory conditions: case report and differential diagnosis. *J Emerg Med* 1992;10(2):151-6.
3. Molderez CM, Wouters KB, Bergmans GB, Michiels GK. Umbilical discharge: a review of 22 cases. *Acta Chir Belg* 1995;95(3):166-9.
4. Cullen TS. *Embryology, anatomy, and diseases of the umbilicus: together with diseases of the urachus*. Philadelphia: WB Saunders; 1916. p. 253-7.

A CMAJ Call for Medical Images: Clinical Vistas

Send us your interesting clinical images!

Through scopes and scanners, on film and computer screens, with ultrasonography and microscopy, clinicians capture stunning images of illness and healing. *CMAJ* invites you to share your normally privy visual perspectives on anatomy, pathology, diagnostic procedures and therapeutic techniques. Let colleagues outside your specialty take a close look at the characteristic signs of rare conditions (Kayser-Fleischer rings in Wilson's disease) or the interior marvels of your clinical terrain (colonoscopic view of an adenomatous polyp). We're also interested in images that take a wider angle on the context of care (a recently cord-clamped newborn on a cold steel scale). If you have original, unpublished images that are beautiful or informative, rare or classic, we'd like to include them in *CMAJ's* Clinical Vistas.

Send your images or queries to:



Editorial Fellow • Canadian Medical Association Journal
1867 Alta Vista Drive • Ottawa ON K1G 3Y6 Canada
 or email pubs@cma.ca