

References

1. Mudge M, Hughes LE. Incisional hernia; a 10 year prospective study of incidence and attitudes. *Br J Surg* 1985;72(1):70-1.
2. Israelsson LA, Jonsson T. Incisional hernia after midline laparotomy: a prospective study. *Eur J Surg* 1996;162(2):125-9.
3. Ooster PJ, Gjode P, Mortensen BB, Mortensen PB, Bartholin J, Gottrup F. Randomized comparison of polyglycolic acid and polyglyconate sutures for abdominal fascial closure after laparotomy in patients with suspected impaired wound healing. *Br J Surg* 1995;82:1080-2.
4. Gecin E, Kocak S, Erszcz S. Recurrence after incisional hernia repair: results and risk factors. *Surg Today* 1996;26(8):607-9.
5. Irvin TT, Koffman CG, Duthie HI. Layer closure of laparotomy wounds with absorbable and nonabsorbable suture materials. *Br J Surg* 1976;78:705-7.
6. Corman ML, Veidenheimer MC, Coller JA. Controlled clinical trial of three suture materials for abdominal wall closure after bowel operation. *Am J Surg* 1981;141:510-3.
7. Carlson MA, Condon RE. Polyglyconate (Maxon) versus nylon suture in midline abdominal incision closure: a prospective randomized trial. *Ann Surg* 1995;61:980-3.
8. Wissing J, van Vroonhoven TJ, Schatzenkerk ME, Veen HF, Ponsen RJ, Jeekel J. Fascia closure after midline laparotomy: results of a randomized trial. *Br J Surg* 1987;74:738-41.
9. Hodgson NC, Malthaner RA, Ostbye T. The search for the ideal method of abdominal fascial closure: a meta-analysis. *Ann Surg* 2000;231(3):436-42.
10. *Canadian Medical Directory*. Don Mills (ON): Southam Publications; 1998.
11. Dillman DA. *Mail and telephone surveys: the total design method*. New York: John Wiley & Sons; 1978.
12. DesCoteaux JG, Sutherland F. Inguinal hernia repair: a survey of Canadian practice patterns. *Can J Surg* 1999;42(2):127-32.
13. Schecter S, Mulvey J, Eisenstat TE. Management of uncomplicated acute diverticulitis: results of a survey. *Dis Colon Rectum* 1999;42(4):470-5.

SESAP Questions  
Questions SESAP

Category 16, Items 11, 12 and 13

11. A patient sustains a facial injury. The computed tomographic (CT) scans are shown. They are diagnostic of a(n)

- (A) orbital “blow-out” fracture
- (B) malar complex fracture
- (C) Le Fort II fracture
- (D) naso-orbital-ethmoid (NOE) fracture
- (E) frontal sinus fracture


12. Clinical findings associated with this injury include all of the following EXCEPT

- (A) subconjunctival hemorrhage
- (B) infraorbital nerve numbness
- (C) flattening of the cheek prominence
- (D) enophthalmos
- (E) lengthening of the face

13. Which of the following findings is an indication for surgical treatment of this injury?

- (A) Periorbital ecchymosis
- (B) Crepitus in the buccal vestibule
- (C) Unilateral epistaxis
- (D) Impaired extraocular muscle function
- (E) Paresthesias of the cheek and hip

For the 2 incomplete statements and 1 question above, select the answer that is best of the 5 given for each item. For the critique of Items 11, 12 and 13, see page 382.



(Reproduced by permission from *SESAP No. 10 1999–2001 Syllabus* Volume 2. For enrolment in the Surgical Education and Self-Assessment Program, please apply to the American College of Surgeons, 633 North St. Clair St., Chicago IL 60611, USA.)