

- microscopic quantification of collagen fibril diameters in the rabbit medial collateral ligament: a baseline for comparison. *Connect Tissue Res* 1989;19:11-25.
32. Frank C, McDonald D, Bray D, Bray R, Rangayyan R, Chimich D, et al. Collagen fibril diameters in the healing adult rabbit medial collateral ligament. *Connect Tissue Res* 1992;27:251-63.
33. Karnovsky M. A formaldehyde-glutaraldehyde fixative of high osmolality for use in electron microscopy. *J Cell Biol* 1965;27: 137A-138A.
34. Venable J, Coggeshell R. A simplified lead citrate stain for use in electron microscopy. *J Cell Biol* 1965;25:407-8.
35. Shrive N, Chimich D, Marchuk L, Wilson J, Brant R, Frank C. Soft-tissue "flaws" are associated with the material properties of the healing rabbit medial collateral liga-
ment. *J Orthop Res* 1995;13:923-9.
36. Spindler KP, Dawson J, Stahlman GC, Davidson JM. Collagen synthesis and biomechanical response to TGF- β 2 in the healing rabbit MCL [abstract]. *Trans Orthop Res Soc* 1996;21:793.
37. Stahlman, G, Spindler K, Dawson J, Davidson J. Do the growth factors TGF- β 2 and PDGF-AB affect the structural properties of the healing medial collateral ligament? [abstract]. *Transactions of the 20th annual meeting of American Orthopaedic Society for Sports Medicine*. Atlanta: 1994; p. 121-2.
38. Lee J, Chamberlin TA, Schreck PJ, Amiel D. In situ localization of growth factors during the early healing of knee ligament [abstract]. *Trans Orthop Res Soc* 1995;20:158.
39. Natsu-ume T, Nakamura N, Shino K, Toritsuka Y, Horibe, S, Ochi T. Temporal and spatial expression of transforming growth factor- β in the healing patellar ligament of the rat. *J Orthop Res* 1997; 15:837-43.
40. Frank S, Madlener M, Warner S. Transforming growth factors β 1, β 2 and β 3 and their receptors are differentially regulated during normal and impaired wound healing. *J Biol Chem* 1996;271:10188-93.
41. Sciore P, Boykiw R, Hart DA. Semiquantitative reverse transcription-polymerase chain reaction analysis of mRNA for growth factors and growth factor receptors from normal and healing rabbit medial collateral ligament tissue. *J Orthop Res* 1998;16:429-37.
42. Bornstein EF, Harisiadis L, Salomon G, Norton J, Sollberg S, Uitto J, et al. Transforming growth factor- β improves healing of radiation-impaired wounds. *J Invest Dermatol* 1991;97:430-4.

SESAP Questions Questions SESAP

Category 11, Items 33 to 35

33. Reliable screening technique
34. Require(s) mediastinal abnormalities to detect significant injury
35. Equivocal findings mandate aortography

For each of the numbered items above, select the applicable lettered word or phrase below

- (A) Helical computed tomography for traumatic aortic disruption
(B) Chest x-ray for traumatic aortic disruption
(C) Both
(D) Neither

For the answers and a critique of items 33 to 35 see page 362.

(Reproduced by permission from *SESAP No. 11 2002-2004 Syllabus*, Volume 2. For enrolment in the Surgical Education and Self-Assessment Program, please apply to the American College of Surgeons, 633 North St. Clair St., Chicago IL 60611-3211, USA; tel 312 202-5000; fax 312 202-5001; email postmaster@facs.org)