

Mesenteric paraganglioma

Suhail Muzaffar, MB BS;* Saira Fatima, MB BS;* Mohammad S. Siddiqui, MB BS;* Naila Kayani, MD;*
Shahid Pervez, PhD;* Asad J. Raja, FRCS†

Paragangliomas are uncommon tumours that have been described in virtually every site in which normal paraganglia are known to occur. Paragangliomas of the carotid body represent the most common group of extra-adrenal paragangliomas. The other less common sites include jugulotympanic, vagal, mediastinal and retroperitoneal. Paragangliomas have been described in the gastrointestinal tract especially in the duodenum. Paraganglioma as a mesenteric mass is extremely rare and only occasional reports have been published.¹⁻³ Our case of mesenteric paraganglioma in a 76-year-old woman is probably only the fourth published case.

Case report

A 76-year-old woman, known to have diabetes mellitus and osteoarthritis, presented with a painless mass of 2 years' duration, which was gradually increasing in size. Examination of abdomen revealed a well-circumscribed mass, measuring 20 cm × 15 cm, that was firm, nontender and mobile. General physical examination and the rest of the systemic examination revealed no other abnormality. Ultrasonography and computed tomography showed a solid, cystic mass with increased vascularity peripherally that appeared to arise from the small-bowel mesentery. Exploratory laparotomy confirmed the presence of a large, solid, cystic mass in the mesentery of the small bowel. The mass was excised. The patient was well without evidence of recurrence or metastasis at 15-month follow-up.

Grossly, the specimen consisted of a round, multilobulated cystic mass filled with blood clots. At the periphery, the lesion was solid and fleshy. Microscopic sections revealed a cellular neoplasm composed of nests and groups of tumour cells separated by fibrovascular connective tissue, giving a characteristic "zellballen" nested pattern. The tumour cells showed monomorphic vesicular nuclei without significant cytologic atypia. Cytoplasm was moderate and granular. Occasional mitotic figures were identified (Fig. 1). Immunohistochemical studies revealed the presence of chromogranin A, synaptophysin and S-100 protein and

the absence of cytokeratins, MNF 116 and anticytokeratin Cam 5.2. On the basis of histologic and immunohistochemical features, a diagnosis of mesenteric paraganglioma was made.

Discussion

Extra-adrenal paragangliomas are rare neoplasms. They may develop anywhere from the neck to the pelvis; however, mostly they arise intra-abdominally.⁴ Generally, the tumours develop in the third to fifth decade of life.⁵ Increased catecholamine secretion is responsible for presenting symptoms in at




FIG. 1. Photomicrograph of the mesenteric mass. Note the typical "zellballen" arrangement of tumour cells (hematoxylin–eosin stain, original magnification × 200).

From the *Department of Pathology and the †Department of Surgery, Aga Khan University Hospital, Karachi, Pakistan

Accepted for publication May 22, 2001.

Correspondence to: Dr. Suhail Muzaffar, Assistant Professor, Department of Pathology, The Aga Khan University Hospital, Stadium Rd., PO Box 3500, Karachi 74800, Pakistan; fax 92 21 493-4294; suhail.muzaffar@aku.edu

least 25% of the cases.⁶ These extra-adrenal paragangliomas tend to behave more aggressively than others, and the incidence of malignant change has been reportedly ranged from 14%⁷ to 50%.⁶ In the present case, the paraganglioma presented as a gradually increasing abdominal mass of 2 years' duration that did not produce any systemic or local obstructive symptoms. The case probably represents the fourth report of this condition in literature.

The recognition of this tumour as a cause of an abdominal mass is extremely important both for surgeons and pathologists. Preoperative image-guided fine-needle aspiration or intraoperative consultation may help an operating surgeon to plan the surgical approach. This lesion

could be a true diagnostic dilemma for a pathologist; however, cytologic appreciation of neuroendocrine features of the tumour cells may help to avoid an erroneous diagnosis of carcinoma or sarcoma.

References

1. el Allame L, Belkhat S, Chkoff L, Mechatt F, Chkoff MR, Benchkroun BA, et al. [Spontaneous hemoperitoneum complicating intraperitoneal paraganglioma.] *J Chir (Paris)* 1989;126:242-7.
2. Badalament RA, Kenworthy P, Pellegrini A, Drago JR. Paraganglioma of urethra. *Urology* 1991;38:76-8.
3. Barnardo DE, Stavrou M, Bourne R,

Bogomoletz W. Primary carcinoid tumour of mesentery. *Hum Pathol* 1984;15:796-8.

4. Carmichael JD, Daniel WA, Lamon EW. Mesenteric chemodectoma. Report of a case. *Arch Surg* 1970;101:630-1.
5. Samaan NA, Hickey RC. Pheochromocytoma. *Semin Oncol* 1987;14:297-305.
6. Sclafani LM, Woodruff JM, Brennan MF. Extraadrenal retroperitoneal paragangliomas: natural history and response to treatment. *Surgery* 1990;108:1124-30.
7. Hayes WS, Davidson AJ, Grimley PM, Hartman DS. Extraadrenal retroperitoneal paraganglioma: clinical, pathologic, and CT findings. *AJR Am J Roentgenol* 1990;155:1247-50.

Un corédacteur pour le JCC

Le *Journal canadien de chirurgie (JCC)* cherche un corédacteur qui jouera un rôle de leadership dans l'évolution continue du Journal.

Le *JCC*, fondé en 1957, est un journal MEDLINE et paraît six fois l'an. L'Association médicale canadienne (AMC) est l'éditeur propriétaire de ce journal parrainé par six sociétés de spécialité.

En collaboration avec un corédacteur, le(la) candidat(e) choisi(e) appuiera le mécanisme d'examen critique par les pairs, sollicitera des articles de grande qualité, prendra des décisions sur l'acceptation ou le rejet de manuscrits, rédigera et sollicitera des éditoriaux, collaborera avec l'éditeur, le conseil de rédaction et les sociétés parraines pour formuler la politique de rédaction et créer des sections et des chroniques intéressantes, approuvera le contenu de numéros, participera à choisir la couverture et les points saillants de chaque numéro, à tenir une base de données d'examineurs et à faire circuler les manuscrits, à produire un rapport annuel des activités du journal, à présider les réunions semi-annuelles du Conseil de rédaction et à choisir de nouveaux membres du conseil de rédaction parmi des candidatures

proposées par les sociétés parraines.

Les candidats doivent avoir le titre de Fellow en chirurgie et une connaissance étendue de la médecine clinique. Ils doivent posséder d'excellentes compétences de communication orale et écrite, de préférence en français et en anglais, ainsi que des compétences de lecture critique très poussées, en particulier des publications sur la médecine scientifique clinique. La connaissance de l'épidémiologie clinique est souhaitable, ainsi que l'expérience à titre d'auteur et d'examineur, de même que l'expérience comme rédacteur d'un journal scientifique. L'enthousiasme et les idées neuves sont essentiels.

Ce poste à temps partiel est assorti d'un mandat de cinq ans renouvelable. Le(la) candidat(e) choisi(e) pourra travailler à partir de son bureau actuel et touchera des honoraires. Publications AMC prend en charge toute la production imprimée et électronique.

Il faut poser sa candidature au plus tard le 20 janvier 2003. Veuillez faire parvenir votre demande et votre curriculum vitae au D^r John Hoey, rédacteur en chef, Publications AMC, Journal canadien de chirurgie, à l'adresse john.hoey@cma.ca