

# Plasticity of peptidergic innervation in healing rabbit medial collateral ligament

Paul T. Salo, MD; Jasmine A. Beye, MSc; Ruth A. Seerattan, BSc; Catherine A. Leonard, MSc; Tyler J. Ivie, BSc, LLB; Robert C. Bray, MD

**Background:** Denervation substantially impairs healing of the medial collateral ligament (MCL). Because normal ligaments are sparsely innervated, we hypothesized that neuropeptide-containing neurons would sprout or proliferate after ligament transection, followed by later regression with healing, in a manner analogous to blood vessels. **Methods:** We transected the right MCL in 9 mature female New Zealand white rabbits and killed 3 rabbits at 2, 6 or 14 weeks. Alternate sets of 12-mm serial sections of healing MCL scars were examined by fluorescent immunohistochemistry for substance P (SP), calcitonin gene-related peptide (CGRP), neuropeptide Y (NPY) and pan-neuronal marker PGP9.5. **Results:** Normal MCLs had few peptidergic fibres located in the epiligament in a perivascular pattern. At 2 weeks, PGP9.5-, SP- and CGRP-positive fibres had increased in the epiligament adjacent to the injury. By 6 weeks, there were increases in CGRP- and PGP9.5-positive fibres in epiligament and scar, with similar but less marked increases in SP-positive fibres. At 14 weeks, there was notable regression of immunostained peptidergic nerve fibres in the scar. **Conclusion:** This experiment shows evidence for a remarkable plasticity of ligament innervation after injury, supporting the idea that neuronal factors play a fundamental role in wound healing.

**Contexte :** La dénervation nuit considérablement à la guérison du ligament latéral interne (LLI). Parce que les ligaments normaux sont peu innervés, nous avons posé en hypothèse que des neurones contenant des neuropeptides feraient leur apparition ou se multiplieraient après une dissection transversale du ligament suivie d'une régression ultérieure avec la guérison, comme le font les vaisseaux sanguins. **Méthodes :** Nous avons procédé à une dissection transversale du LLI droit chez 9 lapines blanches à maturité de la Nouvelle-Zélande et nous en avons tué 3 à 2, 6 ou 14 semaines. On a examiné par immunohistochimie fluorescente des séries alternantes de coupes de 12 mm de cicatrices LLI en guérison pour repérer la substance P (SP), le peptide lié au gène de la calcitonine (*calcitonin gene-related peptide* ou CGRP), le neuropeptide Y (NPY) et le marqueur pan-neuronal PGP9.5. **Résultats :** Les LLI normaux contenaient peu de fibres peptidergiques situées dans l'épiligament disposées de façon périvasculaire. À 2 semaines, les fibres positives pour PGP9.5, SP et CGRP avaient augmenté dans l'épiligament adjacent au traumatisme. À 6 semaines, on a constaté des augmentations des fibres positives pour CGRP et PGP9.5 dans l'épiligament et la cicatrice et des augmentations semblables mais moins marquées des fibres positives pour la SP. À 14 semaines, on a constaté une régression notable des fibres nerveuses peptidergiques immunocolorées dans la cicatrice. **Conclusion :** Cette expérience démontre une plasticité remarquable de l'innervation ligamentaire après une lésion, ce qui appuie le concept selon lequel des facteurs neuronaux jouent un rôle fondamental dans la guérison de la plaie.

Wound healing is a continuous process that can be said to occur in 3 somewhat arbitrary and overlapping stages: initial inflammation followed by proliferation of cells and matrix and then a prolonged remodelling phase.<sup>1</sup> Within hours of an

initial injury, hemorrhage and clot formation occur, accompanied by an inflammatory response that subsequently leads to angiogenesis, cellular proliferation and scar formation. Then, in the following weeks to months, tissue remodelling takes

place. It is reasonable to suggest that any factors promoting inflammation or cellular proliferation are likely to have a significant effect on the outcome of wound healing.

Numerous studies have indicated a potential role for neuropeptides

McCaig Institute for Bone and Joint Health, Department of Surgery, University of Calgary, Calgary, Alta.

Accepted for publication Jan. 22, 2007

**Correspondence to:** Dr. P.T. Salo, Department of Surgery, Faculty of Medicine, University of Calgary, 3330 Hospital Dr. NW, Calgary AB T2N 4N1; fax 403 270-0617; salo@ucalgary.ca

(small peptide-signalling molecules derived from sensory and sympathetic neurons) in wound healing.<sup>2</sup> Substance P (SP) and calcitonin gene-related peptide (CGRP) are potent vasodilators, making these substances particularly important during the initial inflammatory stage of healing.<sup>2</sup> Neuropeptide Y (NPY) is produced by sympathetic neurons, has vasoactive properties and, in concert with CGRP, modulates endothelial and fibroblast cell proliferation associated with angiogenesis.<sup>3,4</sup> Mice lacking the NPY Y2 receptor exhibit impaired skin wound healing.<sup>4</sup>

Other investigations have demonstrated the benefits of an intact peripheral nervous system on wound healing. Most of these studies have focused on well-innervated structures such as skin. In a rat study, denervation via chemical sympathectomy resulted in decreased healing of skin incisions, compared with control animals.<sup>5</sup> Another study in rats revealed impaired fracture healing after peripheral nerve transection.<sup>6</sup> Intact innervation also plays a role in fetal skin wound repair. Stelnicki and colleagues<sup>7</sup> found that innervation was necessary for scarless wound healing in fetal lambs, and in some cases, healing did not occur at all in denervated tissues.

We have previously reported that denervation significantly impairs healing of the rabbit medial collateral ligament (MCL).<sup>8</sup> Six weeks after injury, denervated ligament scars had one-third the blood flow, exhibited two-thirds the angiogenesis response and were 33% weaker when compared with innervated ligament scars. Normal ligament is rather sparsely innervated, with most fibres found in a perivascular distribution in the enveloping epiligament and only rare unmyelinated fibres found within the deep substance of the ligament proper.<sup>9</sup> The unexpectedly profound effect of denervation on the healing of such a poorly innervated tissue suggested that significant changes in the pattern and den-

sity of innervation could be occurring during normal ligament healing.

In the brain and spinal cord, neuronal plasticity is a phenomenon well-known as a substrate for such diverse phenomena as motor learning, memory consolidation and the recovery of function after brain or spinal cord injury<sup>10,11</sup>; however, less attention has been paid to the structural plasticity of sensory and sympathetic terminal arborizations in the peripheral nervous system. Injury induces dramatic changes in ligament vascularity and blood flow,<sup>12,13</sup> but no previous study has investigated the response of the neural elements in ligament at multiple time points after injury.

We hypothesized that after ligament injury the response of neuropeptide-producing (peptidergic) neurons would parallel or presage the vascular response, with initial sprouting and proliferation followed by remodelling and/or regression at later time points. To investigate this intriguing possibility, we elected to examine injured ligaments with immunohistochemistry to identify changes in the pattern and number of peptidergic nerve endings at different times after injury.

## Methods

### *Surgical procedures*

We used 9 skeletally mature female New Zealand white rabbits in this experiment. All animals were given an intravenous injection of 25 mg/mL, 0.16 mL acepromazine maleate (Atravet; Ayerst Veterinary Laboratories) followed by the induction of general inhalational anesthesia (2%–5% halothane and nitrous oxide; 1 L/min O<sub>2</sub>). Using a sterile technique, we completely transected the right MCL of each rabbit. Postoperatively, the animals were given a subcutaneous injection (3 mL) of the antibiotic Duplocillin LA (Intervet), and the injury site was treated with gentamicin antibiotic spray (Gentocin; Schering-Plough

Animal Health). Three rabbits were killed at each of 2, 6 and 14 weeks. Each animal was handled in accordance with the guidelines of the Canadian Council on Animal Care. The University of Calgary Animal Care Committee approved all research protocols. All possible efforts were taken to minimize animal suffering and the number of animals used.

### *MCL transection*

The MCL was approached through a medial incision followed by blunt dissection of overlying connective tissues. A 5-mm spatula was gently inserted under the entire ligament, and the ligament was sharply transected with a scalpel. The cut ends of the ligament were noted to retract spontaneously, leaving a 3-mm gap. The skin was closed with 4–0 nylon sutures.

### *Sample collection*

Animals were killed by an intravenous overdose of sodium pentobarbital (Euthanyl; Bimeda-MTC Animal Health Inc.). The whole MCL was taken from the right leg and immersed in phosphate-buffered 4% paraformaldehyde overnight. The tissue was then cryoprotected in 2 changes of 30% sucrose phosphate buffer overnight, embedded in Tissue-Tek OCT compound (Sakura Finetek) and snap frozen in 2-methyl butane cooled in liquid nitrogen.

### *Histology*

Blocks were stored at –80°C until they were cut into 12 µm sections with a Leica CM3050 cryostat (Leica Microsystems). For each ligament, serial sagittally oriented longitudinal sections were cut through the entire ligament and scar. The sections were alternately collected into 4 sets so that each set comprised every fourth section of the entire series. This allowed us to use 4 different antibodies for each ligament. Sections were

thaw-mounted onto gelatin-coated slides and allowed to dry overnight.

**Immunohistochemistry**

Staining was done in batches that included specimens from the 3 different postinjury time points, all stained at the same time so that inherent variability in the preparation and staining methods would not affect interpretation of the results. Alternate sets of sections were processed for fluorescent immunohistochemistry with polyclonal antibodies specific for SP (Bachem), CGRP (Bachem), NPY (Millipore) or protein gene product 9.5 (PGP9.5) (Serotec). SP and CGRP are peptides found in unmyelinated sensory neurons. Neuropeptide Y is a peptide widely distributed in the brain but restricted to sympathetic nerve fibres in the peripheral nervous system. PGP9.5 is a peptide marker found in virtually all neurons of the peripheral nervous system. Specific antibody staining was visualized by the appropriate Cy3-conjugated fluorescent secondary antibody (Jackson Immunoresearch Laboratories).


Sections were washed (3 × 5 min) in 0.1 M phosphate buffer saline solution/0.1% Triton X-100 (Sigma-Aldrich) and then blocked with the appropriate normal blocking serum at room temperature for 30 minutes. Primary incubation was performed overnight in a humidity chamber at 4°C, with the recommended titres for each antibody being used: PGP9.5, 1:200 in 2% normal rabbit serum; SP, 1:200 in 2% normal goat serum; CGRP, 1:200 in 2% normal goat serum; and NPY, 1:1000 in 2% normal rabbit serum. One or 2 sections from each series were stained identically, with the exception that the primary antibody was omitted to test for nonspecific staining.

The following day the sections were rinsed (3 × 5 min) in PBS and then incubated with the specific Cy3-conjugated secondary antibodies for 1 hour at room temperature

in a humidity chamber. Finally, the sections were washed (3 × 5 min) in PBS, cover slipped with Fluorosave reagent (Calbiochem) and stored in the dark at 4°C.

**Semiquantitative analysis**

We examined the complete set of serial alternate sections for each ligament (40–50 sections for each neuropeptide or marker), which allowed us to assess the presence throughout the tissue of nerve fibres (profiles) containing SP, CGRP, NPY and PGP9.5. We observed slides on a Leitz DMRB fluorescent scope (Leica), using filters appropriate for Cy3 (excitation 515–560 nm). It was evident that changes in the number of nerve fibre profiles were occurring in the scar region, so we limited the histologic examination to that region of the injured ligaments. The scar region was easily distinguished from native ligament tissue by the abrupt transitions from the regular bundles of collagen fibres seen in normal ligament to the amorphous-appearing matrix and increased cellularity of the scar filling the gap left after the injury (Fig. 1). We graded innervation density semiquantitatively as follows: We counted the number of axonal profiles in the scar region by systematically examining every section of the



**FIG. 1.** Low-power view of a longitudinal section of rabbit medial collateral ligament at 6 weeks postinjury, showing the transition zone between normal ligament matrix and scar. Note the organized parallel bundles of crimped collagen fibres in the normal matrix at left (arrow) and the disorganized non-crimped collagen fibres in the scar at right (arrowhead) (hematoxylin-eosin stain, scale bar = 100 µm).

series for each marker. We defined an axonal profile as a homogeneously stained linear or branching sinuous structure with a minimal length of 5 µm. We defined a growth cone (the growing tip of a sprouting neuron) as a similar structure with an amoeboid-appearing tip. Axonal profiles and growth cones could be readily distinguished from background staining, which occurred in an evenly distributed, amorphous or finely granular pattern. We determined the total number of profiles per ligament scar region for each marker. The data are presented in Table 1.

**Results**

**Immunohistochemistry**

Normal MCL was found to have few peptidergic fibres located almost ex-

**Table 1**

**Total numbers of specific peptidergic neuronal profiles in the scar formed at different times after MCL transection\***

Peptide	Rabbit group	Time point, w		
		2	6	14
PGP9.5	A	0	25	39
	B	220	580	10
	C	144	293	111
	Mean	121	299	53
SP	A	0	7	2
	B	140	320	14
	C	21	500	122
	Mean	53	276	46
CGRP	A	25	5	1
	B	40	280	4
	C	52	424	383
	Mean	39	236	133
NPY	A	—	5	4
	B	21	10	0
	C	119	150	43
	Mean	70	55	18

CGRP = calcitonin gene-related peptide; MCL = medial collateral ligament; NPY = neuropeptide Y; PGP9.5 = protein gene product 9.5; SP = substance P; — = no data.


\*Three rabbits were killed at each time point. Staining was carried out in batches (A, B, C). Specimens in the same row were processed together. Consistently poor staining was seen in batch A because of excessive background staining caused by the use of an improperly stored blocking serum.

clusively in the epiligament on the surface of the ligament in a predominantly perivascular pattern. Rarely were fibres seen within the ligament substance proper. After ligament injury, different temporal patterns of staining distribution were seen for each peptidergic marker (summarized in Table 1). We found that the increases in number of nerve fibres were almost exclusively occurring in or near the scar tissue that formed in the gap between the cut ends of the transected ligament. By definition,

scar tissue is formed *de novo* in the gap created by the injury. Therefore, all nerve fibres found in the scar were newly derived by sprouting and ingrowth from preexisting fibres in the surrounding tissue. Bright-field examination of the ligament sections under low power permitted a visual identification of the scar region that was easily distinguished by its increased cellularity and disorganized collagen matrix (Fig. 1).

At 2 weeks, there were moderate increases in the number of profiles

staining for the sensory neuropeptides CGRP and SP (Fig. 2A and 2D respectively), occurring primarily in epiligament adjacent to the developing scar. By 6 weeks postinjury, there were marked increases in the counts of CGRP- (Fig. 2B) and PGP9.5-immunoreactive neurons (Fig. 2H) in both epiligament and scar, with similar but less marked increases in SP labelling (Fig. 2E). All the neuronal markers identified nerve fibres closely associated with blood vessels in the scar and epiligament. CGRP-



**FIG. 2.** Photomontage of typical high-power images of immunohistochemically stained nerve fibres (arrows) at different time points after injury. (A) At 2 weeks, CGRP fibres appear very fine in the early scar. (B) At 6 weeks, CGRP-immunoreactive growth cone-like structures (large arrow head) and sprouting appear, as well as a predominantly perivascular distribution of fibres. (C) At 14 weeks postinjury, there are fewer CGRP-immunoreactive fibres in the scar, with most found again in the epiligament. (D) All SP-positive profiles were found to be perivascular at 2 weeks after injury. (E, F) Similar to the appearance at 2 weeks, SP-positive profiles were only found associated with proliferating vessels within the scar at later time points. (G) Typical image of NPY-containing fibres associated with small arterial vessels in a 6-week postinjury scar. (H) At 6 weeks postinjury, staining for the marker PGP9.5, which stains all nerve fibres, revealed a pattern that combined aspects of the other markers, with most fibres in a perivascular location, and some free in the scar matrix (scale bar = 20 microns). CGRP = calcitonin gene-related peptide; NPY = neuropeptide Y; PGP9.5 = protein gene product 9.5; SP = substance P; v = vessel lumen.

and PGP9.5-immunoreactive fibres were also seen as free nerve endings in the scar matrix. At 14 weeks, there was a regression of immunostained nerve fibres (Fig. 2C and 2F) to numbers slightly greater than those seen in normal controls (not shown). Of note, there were no noticeable changes in the number of NPY-positive fibres at any time point.

## Discussion

We have found evidence of a remarkable plasticity of innervation after ligament injury. The initial sparse distribution of fibres restricted to the epiligament begins to proliferate by 2 weeks and infiltrates the scar by 6 weeks. The number and distribution of nerve terminals then regresses by 14 weeks. Nerve growth into the ligament scar appears to mimic the angiogenic response to injury, which begins in the epiligament, where the greatest number of vessels are found, and then spreads to the midsubstance of the scar.<sup>13,14</sup> The similarity between the vascular and neuronal responses to ligament injury is consistent with recent evidence that growing neurons and vessels are subject to similar, and in some instances identical, regulatory signalling pathways (see Carmeliet<sup>15</sup> for a review).

Some CGRP- and SP-containing profiles were noted in the scar region as early as the 2-week time point, consistent with the known role of these neuropeptides in promoting inflammation. The expression of PGP9.5, a general nerve marker, paralleled both SP and CGRP neuropeptides at the 2-week time point.

Six weeks after ligament injury, a time when cellular proliferation and new matrix production are maximal, we found that there were numerous SP- and CGRP-labelled nerve fibres in the scar. CGRP and PGP9.5 immunolabelling increased more than SP at the 6-week time point. This likely reflects the fact that CGRP-containing sensory neurons are almost 3 times more prevalent in

articular tissues than SP-containing neurons.<sup>16</sup> The peak number of labelled profiles for all the examined neuropeptides was at 6 weeks postinjury, which also coincides with the time of peak blood flow and vascular volume in rabbit MCL.<sup>13,14</sup> NPY staining in the scars did not change significantly after injury. Because NPY is a marker for sympathetic neurons, whereas SP and CGRP are found in sensory neurons, this observation could indicate that sensory and sympathetic neurons respond differently to ligament injury in this model.

At 14 weeks, there were fewer peptidergic profiles in the scars, consistent with our hypothesis that neuronal responses would parallel the angiogenesis response. The dynamic and highly plastic nature of the neuronal response after ligament injury further supports the idea that innervation plays an important role in promoting the early healing response of these tissues. Our findings are remarkably similar to those recently reported for injured Achilles tendon in rats,<sup>17,18</sup> supporting the speculation that neuronal proliferation and activity may be a necessary aspect of wound healing in many tissues and species.

Owing to the small number of animals studied at each time point, the morphological nature of the data, and the amount of interanimal variation, a statistical analysis was not feasible in this study. However, this preliminary study provides clear supporting evidence that an intact peripheral nervous system is dynamically responsive to the early stages of ligament injury. Nerve growth, blood flow and angiogenesis are all substantially upregulated from 2 to 6 weeks after ligament rupture, indicating that these 3 processes likely all influence the healing process in an interdependent manner.

Pain after joint injury is common, severe and disabling, but little is known about the processes and mediators that regulate the activity of the peripheral nervous system in this context. Unlike the phenomenon of

angiogenesis, which has received considerable attention in recent years, the factors involved in neuronal sprouting and growth have received little attention outside the realm of direct injury to brain, spinal cord or peripheral nerve. Further research may reveal the molecules and mechanisms regulating the plasticity of neuronal morphology in injured connective tissue. This could lead to important insights and perhaps new avenues for the treatment of these very common, frequently disabling injuries.

**Acknowledgements:** This work was supported by an operating grant from the Arthritis Society of Canada. Dr. Salo is a Senior Scholar and Dr. Bray a Scientist of the Alberta Heritage Foundation for Medical Research.

**Competing interests:** None declared.

**Contributors:** Drs. Salo and Bray and Mr. Ivie designed the study. Mses. Seerattan and Leonard and Mr. Ivie acquired the data, which Dr. Salo, Ms. Beye and Mr. Ivie analyzed. Drs. Salo and Bray and Mses. Beye and Seerattan wrote the article, which Drs. Salo and Bray, Ms. Leonard and Mr. Ivie reviewed. All authors gave final approval for publication.

## References

1. Clark RAF. Wound repair: overview and general considerations. In: *Molecular and cellular biology of wound repair*. New York: Plenum Press; 1996. p. 3-50.
2. Brain SD. Sensory neuropeptides: their role in inflammation and wound healing. *Immunopharmacology* 1997;37:133-52.
3. Chronwall BM, Zukowska Z. Neuropeptide Y, ubiquitous and elusive. *Peptides* 2004;25:359-63.
4. Ekstrand AJ, Cao R, Bjorndahl M, et al. Deletion of neuropeptide Y (NPY) 2 receptor in mice results in blockage of NPY-induced angiogenesis and delayed wound healing. *Proc Natl Acad Sci U S A* 2003; 100:6033-8.
5. Kim LR, Whelpdale K, Zurowski M, et al. Sympathetic denervation impairs epidermal healing in cutaneous wounds. *Wound Repair Regen* 1998;6:194-201.
6. Madsen JE, Hukkanen M, Aune AK, et al. Fracture healing and callus innervation after peripheral nerve resection in rats. *Clin Orthop Relat Res* 1998;351:230-40.
7. Stelnicki EJ, Doolabh V, Lee S, et al. Nerve dependency in scarless fetal wound healing. *Plast Reconstr Surg* 2000;105: 140-7.

8. Ivie TJ, Bray RC, Salo PT. Denervation impairs healing of the rabbit medial collateral ligament. *J Orthop Res* 2002;20:990-5.
9. McDougall JJ, Bray RC, Sharkey KA. Morphological and immunohistochemical examination of nerves in normal and injured collateral ligaments of rat, rabbit, and human joints. *Anat Rec* 1997;248:29-39.
10. Rossini PM, Dal Forno G. Neuronal post-stroke plasticity in the adult. *Restor Neurol Neurosci* 2004;22:193-206.
11. Straub VA, Styles BJ, Ireland JS, et al. Central localization of plasticity involved in appetitive conditioning in *Lymnaea*. *Learn Mem* 2004;11:787-93.
12. Bray RC, Butterwick DJ, Doschak MR, et al. Coloured microsphere assessment of blood flow to knee ligaments in adult rabbits: effects of injury. *J Orthop Res* 1996;14:618-25.
13. Bray RC, Leonard CA, Salo PT. Correlation of healing capacity with vascular response in the anterior cruciate and medial collateral ligaments of the rabbit. *J Orthop Res* 2003;21:1118-23.
14. Bray RC, Leonard CA, Salo PT. Vascular physiology and long-term healing of partial ligament tears. *J Orthop Res* 2002;20:984-9.
15. Carmeliet P. Blood vessels and nerves: common signals, pathways and diseases. *Nat Rev Genet* 2003;4:710-20.
16. Salo PT, Theriault ET. Number, distribution and neuropeptide content of rat knee joint afferents. *J Anat* 1997;190:515-22.
17. Ackermann PW, Ahmed M, Kreicbergs A. Early nerve regeneration after Achilles tendon rupture—a prerequisite for healing? A study in the rat. *J Orthop Res* 2002;20:849-56.
18. Ackermann PW, Li J, Lundeberg T, et al. Neuronal plasticity in relation to nociception and healing of rat Achilles tendon. *J Orthop Res* 2003;21:432-41.

## Canadian Surgery FORUM canadien de chirurgie

The Canadian Surgery FORUM canadien de chirurgie will hold its annual meeting from Sept. 11 to 14, 2008, in Halifax, Nova Scotia. This interdisciplinary meeting provides an opportunity for surgeons across Canada with shared interests in clinical practice, continuing professional development, research and medical education to meet in a collegial fashion. The scientific program offers material of interest to academic and community surgeons, residents in training and students.

The major sponsoring organizations include the following:

- The Canadian Association of General Surgeons
- The Canadian Society of Colon and Rectal Surgeons
- The Canadian Association of Thoracic Surgeons
- The Canadian Society of Surgical Oncology

Other participating societies include the American College of Surgeons, the Canadian Association of Surgical Chairmen, the Canadian Association of University Surgeons, the Canadian Hepato-Pancreato-Biliary Society, the Canadian Undergraduate Surgical Education Committee, Doctors Nova Scotia, the James IV Association of Surgeons, the Ontario Association of General Surgeons and the Trauma Association of Canada.

For registration and further information contact [surgeryforum@rcpsc.edu](mailto:surgeryforum@rcpsc.edu); [www.cags-accg.ca/cagsaccg.php?page=56](http://www.cags-accg.ca/cagsaccg.php?page=56)

## Canadian Surgery FORUM canadien de chirurgie

La réunion annuelle du Canadian Surgery FORUM canadien de chirurgie aura lieu du 11 au 14 septembre 2008 à Halifax en Nouvelle Écosse. Cette réunion interdisciplinaire permet aux chirurgiens de toutes les régions du Canada qui s'intéressent à la pratique clinique, au perfectionnement professionnel continu, à la recherche et à l'éducation médicale d'échanger dans un climat de collégialité. Un programme scientifique intéressera les chirurgiens universitaires et communautaires, les résidents en formation et les étudiants.

Les principales organisations qui parrainent cette réunion sont les suivantes :

- L'Association canadienne des chirurgiens généraux
- La Société canadienne des chirurgiens du côlon et du rectum
- La Société canadienne de chirurgie thoracique
- La Société canadienne d'oncologie chirurgicale

Le American College of Surgeons, le Canadian Association of Surgical Chairmen, l'Association canadienne des chirurgiens universitaires, le Canadian Hepato-Pancreato-Biliary Society, le Comité canadien de l'éducation chirurgicale de premier cycle, Doctors Nova Scotia, l'Association des chirurgiens James IV, le Ontario Association of General Surgeons, et l'Association canadienne de traumatologie sont au nombre des sociétés qui appuient cette activité.

Pour vous inscrire, veuillez communiquer à [surgeryforum@rcpsc.edu](mailto:surgeryforum@rcpsc.edu); [www.cags-accg.ca/cagsaccg.php?page=56](http://www.cags-accg.ca/cagsaccg.php?page=56)