

SURGICAL IMAGES: SOFT TISSUE

Mucinous cystadenoma of the appendix

Simon S.M. Ng, MBChB (Hons)
Janet F.Y. Lee, MD

From the Department of Surgery,
The Chinese University of Hong Kong,
Prince of Wales Hospital, Shatin, Hong
Kong SAR, China

Correspondence to:

Dr. S.S.M. Ng
Department of Surgery
The Chinese University of Hong Kong
Prince of Wales Hospital
Shatin, Hong Kong SAR
China
fax 852 26377974
simonng@surgery.cuhk.edu.hk

A 61-year-old man presented with chronic intermittent right iliac fossa pain that had lasted for a few years. His medical history included chronic obstructive airway disease and old pulmonary tuberculosis. Colonoscopy revealed a 3-cm smooth bulbous submucosal mass at the cecum just behind the ileocecal valve (Fig. 1). Biopsy of the lesion did not confirm any disease. A contrast-enhanced computed tomography (CT) scan of the abdomen revealed a hypodense and distended tubular structure (1.7×5.3 cm) that originated from the cecal pole, the features of which were compatible with a mucinous neoplasm of the appendix (Fig. 2). Because the base of the appendiceal mass was broad and seemed to be contiguous with the cecal wall on CT, we considered an appendectomy alone to be unsafe because there would be considerable risk of peritoneal spillage of mucin during transection of the appendix. We performed a laparoscopic-assisted right hemicolectomy instead (Fig. 3). Histopathological examination of the resected specimen revealed mucinous cystadenoma of the appendix without evidence of cancer. The patient made an uneventful recovery after surgery and was discharged on postoperative day 7.

Mucinous cystadenoma is a rare cystic neoplasm of the veriform appendix characterized by villous adenomatous changes of the appendiceal epithelium associated with marked distension of the appendiceal lumen with mucin. The most common presentation is right iliac fossa pain, similar to an acute appendicitis; however, about 25% of patients are asymptomatic and the condition is found incidentally on imaging or at the time of surgery.^{1,2} Other reported complications include intestinal obstruction, intussusception, gastrointestinal bleeding and extrinsic ureteral compression. The most fearful complication is pseudomyxoma peritonei secondary to spontaneous or iatrogenic rupture of the appendix and consequent spillage of neoplastic cells and mucin into the peritoneal cavity. A correct preoperative diagnosis is thus important to help in




Fig. 1: Colonoscopic image showing a 3-cm smooth bulbous submucosal mass at the cecum.



Fig. 2: Contrast-enhanced computed tomography scan of the abdomen showing a hypodense and distended tubular structure originating from the cecal pole (arrow).



Fig. 3: The right hemicolecotomy specimen.

the choice of surgical tactics and to avoid iatrogenic rupture and peritoneal spillage of mucin during surgery.

As illustrated in our patient, both colonoscopy and CT can facilitate the preoperative diagnosis of mucinous cystadenoma of the appendix. The use of laparoscopy can allow a better evaluation of the lesion intraoperatively and assist in surgical planning. With the availability of surgeons experienced in minimally invasive surgery, laparoscopic-assisted right hemicolecotomy can be safely performed for mucinous cystadenoma or cystadenocarcinoma of the appendix to minimize the unnecessary complication of peritoneal spillage of mucin, which may occur during open or laparoscopic appendectomy. Moreover, the patients benefit from minimally invasive surgery, including a smaller wound and faster convalescent period.³

Competing interests: None declared.

References

- Rampone B, Roviello F, Marrelli D, et al. Giant appendiceal mucocele: report of a case and brief review. *World J Gastroenterol* 2005;11:4761-3.
- Zanati SA, Martin JA, Baker JP, et al. Colonoscopic diagnosis of mucocele of the appendix. *Gastrointest Endosc* 2005;62:452-6.
- Lau H, Yuen WK, Loong F, et al. Laparoscopic resection of an appendiceal mucocele. *Surg Laparosc Endosc Percutan Tech* 2002;12:367-70.

How you can get involved in the CMA!

The CMA is committed to providing leadership for physicians and promoting the highest standard of health and health care for Canadians. To strengthen the association and be truly representative of all Canadian physicians the CMA needs to hear from members interested in serving in elected positions and on appointed committees and advisory groups. The CMA structure comprises both governing bodies and advisory bodies either elected by General Council or appointed by the CMA Board of Directors. The Board of Directors — elected by General Council — has provincial/territorial, resident and student representation, is responsible for the overall operation of the CMA and reports to General Council on issues of governance.

CMA committees advise the Board of Directors and make recommendations on specific issues of concern to physicians and the public. Five core committees mainly consist of regional, resident and student representation while other statutory and special committees and task forces consist of individuals with interest and expertise in subject-specific fields. Positions on one or more of these committees may become available in the coming year.

For further information on how you can get involved, please contact:

Cherise Araujo, Corporate and Governance Services
Canadian Medical Association
1867 Alta Vista Drive, Ottawa ON K1G 5W8
Fax 613 526-7570, Tel 800 663-7336 x2199
involved@cma.ca

By getting involved, you will have an opportunity to make a difference.

We hope to hear from you!

ASSOCIATION
MÉDICALE
CANADIENNE  CANADIAN
MEDICAL
ASSOCIATION